

Exclusive Compton Scattering on The Proton

J. P. Chen, E. Chudakov, C. DeJager, P. Degtyarenko,
R. Ent, J. Gomez, O. Hansen, C. Keppel, M. Kuss, J. LeRose,
M. Liang, R. Michael, J. Mitchell, S. Nanda, P. Rutt,
A. Saha, B.Wojtsekhowski (co-spokesman and contact person)
TJNAF, Newport News VA

B. S. Bains, M. Bouwhuis, R. J. Holt, N.C.R. Makins,
A. M. Nathan (co-spokesman), S.E. Williamson
University of Illinois at Urbana-Champaign, Urbana IL

G. Dodge, C. Hyde-Wright (co-spokesman),
A. Radyushkin, L. B. Weinstein, P. Ulmer
Old Dominion University, Norfolk VA

P. Bosted
The American University, Washington DC

J. M. Finn
College of William and Mary, Williamsburg VA

R. Gilman
Rutgers University, New Brunswick NJ

J. Berthot, P. Bertin, V. Breton, H. Fonvieille, Y. Roblin
University Blaise Pascal/IN2P3, Aubière FRANCE

W. Bertozzi, K. Fissum, J. Gao, S. Gilad, N. Liyanage,
D. Rowntree, J. Zhao, Z. Zhou
Massachusetts Institute of Technology

G. Chang
University of Maryland

A. Ketikyan, A. Shahinyan, H. Voskanyan
Yerevan Physics Institute

W. Boeglin, P. Markowitz
Florida International University

J. Hines, G. Strobel, J. Templon
University of Georgia

G. Feldman
George Washington University

C. L. Morris
Los Alamos National Laboratory

V. Gladyshev, R. A. Lindgren
University of Virginia

J. Calarco, W. Hersman
University of New Hampshire

A. Gasparian
Hampton University
(December 21, 1997)

Abstract

We propose measurements at 3 – 6 GeV of Compton scattering $\gamma + p \rightarrow \gamma + p$ at large perpendicular momentum transfer to the proton. Recent theoretical developments offer the possibility of a unified treatment of high energy Compton scattering, elastic form factors, and deep inelastic scattering. We will use an untagged bremsstrahlung beam in Hall A with coincidence detection of the recoil proton and scattered photon. The proton will be detected in one of the HRS spectrometers, and the photon will be detected in a large-area, multi-segment Pb-Glass calorimeter.